

Cat Amputation Aftercare Information

Susan Hodge, DVM, DACVS-SA

Date of Surgery: _____

Amputation of: Right Left
 Forelimb Hind limb

Convalescent Period: Most pets begin to adjust to walking on three limbs well and in about 7 to 10 days your pet should be ambulating well. Some pets will experience a few days of mild depression following surgery which can be minimized by providing comfortable bedding, bland food offered by hand feeding, and extra comforting. Please call if your pet does not progress as stated above or you have concerns.

Medications:

- Fentanyl patch** (sizes available include 12, 25, 50, 75, & 100 ug patches): This is a clear plastic patch applied to your pet's skin for relief of discomfort or pain control. You will be instructed on when to remove the patch from your pet's skin (usually the date it should be removed will be noted). In most pets, the effects of this patch will work for about 3-5 days total (average of 4 days). It can be disposed of in two ways only, either place the patch in a plastic bag and return it to the office that dispensed it to you for disposal, or fold it in half, and flush it down the toilet. Do not allow pets or people to lick, eat or chew the patch because this may result in intoxication. The effects of the patch take 12-24 hours to either start to work or for the effects of the patch to wear off. Other medications can be used with the patch in place if needed and instructions on adjusting doses should be provided for you.
- Pain relieving medications: **Buprenex 0.5 mg/mL**: The dose will be given in terms of the number of mL (1 syringe full of pre-filled medication) needed to be administered by mouth and the frequency will be noted for you (either every 8, or 12 hours) for 10-14 days as needed for pain control. This medication is an opioid and may cause sedation, mild constipation, or a decreased appetite. These side effects are related to the dose of the medication. Please call your doctor if you have concerns about any side effects. If a Fentanyl patch has been applied to your pet, this medication may not be needed until the patch has been removed, however it can be started with the Fentanyl patch at half the recommended dose if needed. Please call if you have additional questions or concerns.
 - The first doses are due about 8-12 hours after removal of the Fentanyl patch
 - OR The time of the next dose that this medication is due will be noted on your discharge instructions (_____ am/pm).
- Other: Antibiotics** may be administered if needed, depending on the case. The dose will be given in terms of the number of tablet(s)/capsule(s) needed to be administered by mouth and the frequency will be noted for you (either every 12 or 24 hours) for the appropriate number of days. Antibiotics would be used to prevent infection or treat contamination noted at the time of surgery. Any antibiotic may cause nausea, vomiting, or diarrhea. Most side effects can be minimized by administering the medication with food. The time of the next dose that this medication is due will be noted on your discharge instructions (_____ am/pm).
- Anti-inflammatory medications: **Non-steroidal anti-inflammatory (NSAID's) are not always used in cats, but options include Metacam or Onsiar® (Robenacoxib)** : (tablet sizes or liquid concentrations vary depending on the product): The dose will be given in terms of the number of

tablet(s), partial tablet, or volume of liquid (in mL) needed to be administered by mouth and the frequency will be noted for you (either every 12 or 24 hours) for 10-14 days as needed for pain control. These medications reduce inflammation, pain, and swelling. Do not give any other medications (particularly NOT Aspirin or steroids) with this medicine unless instructed by a veterinarian. Stop administering this medication if you notice any vomiting, diarrhea, loss of appetite, or black tarry stool. The time of the next dose that this medication is due will be noted on your discharge instructions (_____ am/pm). Some doctors prefer that this medication is not started until your pet leaves the hospital and is eating more normally.

Other:

Exercise: Limit activity to a small room or large dog crate for 2-3 weeks. Do not allow your pet to be jumping, climbing stairs, or roughhousing with other pets or people. Strenuous activity can breakdown the surgical repair, cause pain, cause the incision to open, and prevent a successful recovery.

- Short periods of time on a single level can be allowed with direct supervision.
- Please do not let your pet outside during recovery.

LAB /BIOPSY RESULTS: Biopsy results should return in 5-7 days (up to 10 days if bone was submitted). Dr. Hodge will call you with these results, but please call if you have not heard from her in 7 days. A copy of these results will be sent to your regular veterinarian after review.

LITTER PAN – Use shredded paper or Yesterdays News for litter for 10-14 days.

Physical therapy may help your pet recover better. Please contact us with additional questions about physical therapy.

Incision: The incision may be covered by a bandage, however if the bandage slips or needs to be removed, please check the incision daily for signs of problems including redness, swelling, pain, discharge, or loss of sutures/staples. Please call if any of these are noted. Do not allow your pet to lick or scratch at the incision. Please obtain an Elizabethan collar from us, your regular veterinarian, or a pet supply store to prevent problems with the incision. Another option is to place bitter apple solution around the incision or to place a t-shirt on your pet depending on their tolerance of these pieces of clothing. If your pet does not have a bandage, there may be a small band-aid pad covering the incision that can be removed in 1-2 days or sooner if it falls off. It is intended to keep the incision clean while in the hospital.

- There are no external staples/sutures to be removed, but the incision should still be checked.
- There are staples/sutures that need to be removed in about 10-14 days.

BANDAGES: A bandage has been applied to help keep the incision covered for comfort and for cleanliness. The bandage should be kept clean and dry. Placing a t-shirt or sock over the bandage will not only help keep the bandage cleaner, it will allow it to last longer. Please call if the bandage slips, gets wet, soiled, or there is an odor.

- A bandage check/change in 7-10 days.
- A bandage removal in 10-14 days.
- A drain is present. Most drains will be removed in 3-5 days.
- Additional instructions:

Nausea is common following General Anesthesia, in case of nausea:

- **FOOD AND WATER:** Large intakes of food and water following a hospital visit are common due to the excitement of returning home. This type of vomiting can often be eliminated by not allowing access to food or water for 2 hours after returning home.
 - o A bland/special diet may be recommended, the length of time will depend on the type of surgery that has been performed. For orthopedic or neurologic surgery the bland diet may only be needed for a few days, following intestinal surgery, the special diet may be needed for up to a couple of weeks.
 - o Following any nausea or intestinal surgery, food and water is recommended in small amounts every 4-6 hours for a few days, then the volume can be increased and the frequency of meal feedings can be decreased to what is more normal.

A bland diet such as Eukanuba Low Residue, Hill's I/D, or Royal Canine Intestinal diets can be purchased or a home-made diet can be made with a 50:50 mixture of a cooked protein source such as hamburger, turkey, chicken, or cottage cheese (all fat removed) combined with a carbohydrate source such as rice, pasta, or mashed potato. We recommend gradually mixing the regular diet into this bland diet during the first week home.

**** PLEASE RETURN your pet for an incision check (staple removal) in 10-14 days