

Cat Femoral Head & Neck Osteotomy (FHO – Hip) Aftercare Information

Susan Hodge, DVM, DACVS-SA

Surgery Date: _____

Surgery Performed: FHO of () Right () Left hip

Convalescent period: Your pet should be walking and touching the toes of the operated limb to the ground at a walk within 10-14 days. From this point on, the use of the leg should improve and the degree of lameness should gradually resolve. If you notice that your pet worsens or loses the ability to use the limb at any time during recovery, please give us a call.

Prognosis: Most small dogs and cats do very well following the FHO surgery and your pet should walk well. You may notice a change in your pet's gait (how they walk), but the previously noted pain will have resolved. The goal following an FHO surgery is to remove bone which is rubbing and causing pain (sometimes related to a fracture, arthritis, or hip luxation), then to allow the joint to fill in with scar tissue for support. The total healing time take a few weeks, however we want to be sure that the scarring allows enough movement, thus early activity and physical therapy is recommended. By 2 weeks following surgery we recommend the re-initiation of activity to allow stretching of the healing tissue in order to prevent too much loss of motion to the hip joint. Physical therapy exercises can be started (range of motion in particular) as early as a few days following surgery.

Medications:

- () **Fentanyl patch** (sizes available include 12, 25, 50, 75, & 100 ug patches): This is a clear plastic patch applied to your pet's skin for relief of discomfort or pain control. You will be instructed on when to remove the patch from your pet's skin (usually the date it should be removed will be noted). In most pets, the effects of this patch will work for about 3-5 days total (average of 4 days). It can be disposed of in two ways only, either place the patch in a plastic bag and return it to the office that dispensed it to you for disposal, or fold it in half, and flush it down the toilet. Do not allow pets or people to lick, eat or chew the patch because this may result in intoxication. The effects of the patch take 12-24 hours to either start to work or for the effects of the patch to wear off. Other medications can be used with the patch in place if needed and instructions on adjusting doses should be provided for you.
 - () Pain relieving medications: **Buprenex 0.5 mg/mL**: The dose will be given in terms of the number of mL (1 syringe full of pre-filled medication) needed to be administered by mouth and the frequency will be noted for you (either every 8, or 12 hours) for 10-14 days as needed for pain control. This medication is an opioid and may cause sedation, mild constipation, or a decreased appetite. These side effects are related to the dose of the medication. Please call your doctor if you have concerns about any side effects. If a Fentanyl patch has been applied to your pet, this medication may not be needed until the patch has been removed, however it can be started with the Fentanyl patch at half the recommended dose if needed. Please call if you have additional questions or concerns.
 - () The first doses are due about 8-12 hours after removal of the Fentanyl patch
- OR () The time of the next dose that this medication is due will be noted on your discharge instructions (_____ am/pm).

Other: Antibiotics may be administered if needed, depending on the case. The dose will be given in terms of the number of tablet(s)/capsule(s) needed to be administered by mouth and the frequency will be noted for you (either every 12 or 24 hours) for the appropriate number of days. Antibiotics would be used to prevent infection or treat contamination noted at the time of surgery. Any antibiotic may cause nausea, vomiting, or diarrhea. Most side effects can be minimized by administering the medication with food. The time of the next dose that this medication is due will be noted on your discharge instructions (_____ am/pm).

Anti-inflammatory medications: **Non-steroidal anti-inflammatory (NSAID's) are not always used in cats, but options include Metacam or Onsior® (Robenacoxib)** : (tablet sizes or liquid concentrations vary depending on the product): The dose will be given in terms of the number of tablet(s), partial tablet, or volume of liquid (in mL) needed to be administered by mouth and the frequency will be noted for you (either every 12 or 24 hours) for 10-14 days as needed for pain control. These medications reduce inflammation, pain, and swelling. Do not give any other medications (particularly NOT Aspirin or steroids) with this medicine unless instructed by a veterinarian. Stop administering this medication if you notice any vomiting, diarrhea, loss of appetite, or black tarry stool. The time of the next dose that this medication is due will be noted on your discharge instructions (_____ am/pm). Some doctors prefer that this medication is not started until your pet leaves the hospital and is eating more normally.

Other:

Exercise: Limit activity to a large crate or a small room without furniture for 2 weeks (food and water can be left in these areas). Limit outdoor activity to very short leash walks for bowel and urinary purposes only for the next 2 weeks. Do not allow jumping, climbing stairs, or roughhousing with other pets or people for the full length of recovery. Strenuous activity can breakdown the surgical repair, cause pain, cause the incision to open, and prevent a successful recovery.

- Short periods of time on a single level can be allowed with direct supervision.
- Please do not let your pet outside during recovery.

LITTER PAN – Use shredded paper or Yesterdays News instead of regular litter for 14 days. If your pet will not go to the bathroom in this material, you can cover the regular litter box with a plastic bag (and place some of the shredded paper on top of the plastic to absorb urine). This will allow your pet to feel the litter under the bag and they will be more likely to use the box this way.

Incision: Please check the incision daily for signs of problems including: redness, swelling, pain, discharge, loss of sutures, or opening of the incision. Please call our office if you have concerns about the incision. Do not allow your pet to lick at the incision. Please use the Elizabethan collar from us, your regular veterinarian, or a pet supply store to prevent licking or chewing at the incision. Another option is to apply bitter apple solution to the skin around the incision several times a day to prevent licking. Your pet should return to Dr. Hodge in about 2 weeks to check the incision. There may be a small band-aid and non-stick pad covering the incision. This can be removed in 1-2 days or sooner if it falls off. It is intended to keep the incision clean while in the hospital.

- There are no external staples/sutures to be removed, but the incision should still be checked.
- There are staples/sutures that need to be removed in about 10-14 days.

Physical Rehabilitation: Physical therapy is recommended for optimal results.

- Initial therapy: Some swelling and bruising is expected to occur after surgery which may be noted about 2-3 days after surgery, and it may take about 5-7 days for the swelling to fully resolve. You can help minimize swelling in the leg by applying a cold compresses to the incision site (hip) for the first few days. Place a cold pack on the leg 2-3 times daily for 10-15 minute sessions for the first 3 days after surgery if able. A plastic bag of frozen vegetables makes a good cold pack. On day 4, we recommend changing to a warm compress, which can be applied to the limb 2-3 times a day for 10-15 minutes if swelling is still present. **** The goal of the cold or warm therapy is to help relieve both pain and swelling, however if your pet fights or resists this treatment, do not continue as they will recover regardless of your ability to apply compresses to the limb.
 - () Schedule suture/staple removal or an incision check in 10-14 days.
 - () Contact us with additional questions about physical therapy.
 - () Other: Laser therapy can be initiated as soon as following surgery. Laser therapy can be used 2-3 times a week in the initial few weeks following surgery to help with healing if desired.

Nausea is common following General Anesthesia, in case of nausea:

- FOOD AND WATER: Large intakes of food and water following a hospital visit are common due to the excitement of returning home. This type of vomiting can often be eliminated by not allowing access to food or water for 2 hours after returning home.
 - o A bland/special diet may recommended, the length of time will depend on the type of surgery that has been performed. For orthopedic or neurologic surgery the bland diet may only be needed for a few days, following intestinal surgery, the special diet may be needed for up to a couple of weeks.
 - o Following any nausea or intestinal surgery, food and water is recommended in small amounts every 4-6 hours for a few days, then the volume can be increased and the frequency of meal feedings can be decreased to what is more normal.

A bland diet such as Eukanuba Low Residue, Hill's I/D, or Royal Canine Intestinal diets can be purchase or a home-made diet can be made with a 50:50 mixture of a cooked protein source such as hamburger, turkey, chicken, or cottage cheese (all fat removed) combined with a carbohydrate source such as rice, pasta, or mashed potato. We recommend gradually mixing the regular diet into this bland diet during the first week home.

PLEASE RETURN your pet for () An incision check (suture removal) is recommended in 10-14 days
() A re-examination is recommended in 6 weeks