

TPLO with NOCITA Aftercare Information

Susan Hodge, DVM, DACVS-SA

Surgery Date: _____

Surgery Performed: TPLO on () right () left stifle.

Convalescent period: Your pet should be walking and touching the toes of the operated limb to the ground at a walk within 10-14 days. From this point on, the use of the leg should improve and the degree of lameness should gradually resolve. If you notice that your pet worsens or loses the ability to use the limb at any time during recovery, please give us a call.

Prognosis: By 8 weeks after surgery the stability in the knee improves (the cut in the bone has almost healed), and your pet should be using the limb well. After healing takes place, your pet should be able to walk normally on the leg again. During times of weather changes (cold or damp) or if out in the cold weather for a while, some pets will experience some discomfort noted as lameness in the leg.

Medications:

- (X) **NOCITA®:** Your pet got an infusion of this long-acting local medication at the time the incision was closed on _____. This product will release slowly over the next 72 hours to provide constant pain control. You can start the oral pain medication if needed while this product is working, but you may not feel the need for pain medication for 3 days (evening of _____). The injection of this product may cause some bruising at the incision site, but this will resolve without treatment.
- () **Anti-inflammatory medications:** Non-steroidal anti-inflammatory (NSAID's) include Rimadyl (Novox, Carprofen), Deramaxx, Metacam, Previcox, and others are available in either Brand name or generic forms: (tablet sizes or liquid concentrations vary depending on the product): The dose will be given in terms of the number of tablet(s), partial tablet, or volume of liquid (in mL) needed to be administered by mouth and the frequency will be noted for you (either every 12 or 24 hours) for 10-14 days as needed for pain control. These medications reduce inflammation, pain, and swelling. Do not give any other medications (particularly NOT Aspirin or steroids) with this medicine unless instructed by a veterinarian. Stop administering this medication if you notice any vomiting, diarrhea, loss of appetite, or black tarry stool. The time of the next dose that this medication is due will be noted on your discharge instructions (_____ am/pm). Some doctors prefer that this medication is not started until your pet leaves the hospital and is eating more normally.
- () **Pain relieving medications: Tramadol 50 mg:** The dose will be given in terms of the number of tablet(s) or partial tablet needed to be administered by mouth and the frequency will be noted for you (either every 8, or 12 hours) for 10-14 days as needed for pain control. This medication is an opioid and may cause sedation, mild constipation, or a decreased appetite. These side effects are related to the dose of the medication. Please call if you have concerns about any side effects. If your pet was given NOCITA, this medication may not be needed, however it can be started at any time. The time of the next dose that this medication is due will be noted on your discharge instructions (_____ am/pm).
- () **Other: Antibiotics** may be administered if needed, depending on the case. The dose will be given in terms of the number of tablet(s)/capsule(s) needed to be administered by mouth and the frequency will be noted for you (either every 12 or 24 hours) for the appropriate number of days. Antibiotics would be used to prevent infection or treat contamination noted at the time of surgery. Any antibiotic may cause nausea, vomiting, or diarrhea. Most side effects can be minimized by administering the medication with food. The time of the next dose that this medication is due will be noted on your discharge instructions (_____ am/pm).

() Other: Sedatives may be sent home for management of activity.

Exercise: Limit activity to a large crate or a small room without furniture for 8 weeks (food and water can be left in these areas). Limit outdoor activity to very short leash walks for bowel and urinary purposes only for the next 8 weeks. Never leave your pet outside, or take your pet off the leash outside during the recovery period. Do not allow jumping, climbing stairs, or roughhousing with other pets or people for the full length of recovery. Strenuous activity can breakdown the surgical repair, cause pain, cause the incision to open, and prevent a successful recovery.

() Short walks are permitted on a leash only (10 minutes maximum). Do not leave your pet off the leash or unattended outside during the time of exercise restriction.

Incision: Please check the incision daily for signs of problems including: redness, swelling, pain, discharge, loss of sutures, or opening of the incision. Please call our office if you have concerns about the incision. Do not allow your pet to lick at the incision. Please use the Elizabethan collar from us, your regular veterinarian, or a pet supply store to prevent licking or chewing at the incision. Another option is to apply bitter apple solution to the skin around the incision several times a day to prevent licking. Your pet should return to Dr. Hodge in about 2 weeks to check the incision. There may be a small band-aid and non-stick pad covering the incision. This can be removed in 1-2 days or sooner if it falls off. It is intended to keep the incision clean while in the hospital.

() There are no external staples/sutures to be removed, but the incision should still be checked.

() There are staples/sutures that need to be removed in about 10-14 days.

Physical Rehabilitation: Physical therapy is recommended for optimal results.

- Initial therapy: Some swelling and bruising is expected to occur after surgery which may be noted about 2-3 days after surgery, especially at the knee and/or at the ankle joint, and it may take about 5-7 days for the swelling to fully resolve. You can help minimize swelling in the leg by applying a cold compresses to the incision site (leg) for the first few days. Place a cold pack on the leg 2-3 times daily for 10-15 minute sessions for the first 3 days after surgery if able. A plastic bag of frozen vegetables makes a good cold pack. On day 4, we recommend changing to a warm compress, which can be applied to the limb 2-3 times a day for 10-15 minutes if swelling is still present. **** The goal of the cold or warm therapy is to help relieve both pain and swelling, however if your pet fights or resists this treatment, do not continue as they will recover regardless of your ability to apply compresses to the limb.

() Schedule suture/staple removal or an incision check in 10-14 days.

() Contact us with additional questions about physical therapy.

() Other: Laser therapy can be initiated as soon as following surgery. Laser therapy can be used 2-3 times a week in the initial few weeks following surgery to help with healing if desired.

Nausea is common following General Anesthesia, in case of nausea:

- **FOOD AND WATER:** Large intakes of food and water following a hospital visit are common due to the excitement of returning home. This type of vomiting can often be eliminated by not allowing access to food or water for 2 hours after returning home.
 - o A bland/special diet may recommended, the length of time will depend on the type of surgery that has been performed. For orthopedic or neurologic surgery the bland diet may only be needed for a few days, following intestinal surgery, the special diet may be needed for up to a couple of weeks.
 - o Following any nausea or intestinal surgery, food and water is recommended in small amounts every 4-6 hours for a few days, then the volume can be increased and the frequency of meal feedings can be decreased to what is more normal.

A bland diet such as Eukanuba Low Residue, Hill's I/D, or Royal Canine Intestinal diets can be purchase or a home-made diet can be made with a 50:50 mixture of a cooked protein source such as hamburger, turkey, chicken, or cottage cheese (all fat removed) combined with a carbohydrate source such as rice, pasta, or mashed potato. We recommend gradually mixing the regular diet into this bland diet during the first week home.

PLEASE RETURN your pet for An incision check (suture removal) is recommended in 10-14 days
 A re-examination is recommended in 8-9 weeks
 A follow-up X-ray is recommended in 8-9 weeks

Additional Information for the Future: Following TPLO surgery, the implants (bone plate) applied to the bone may need to be removed in the future (either because it loosens, becomes exposed through the skin, becomes infected, or causes irritation). Removal of the bone plate is an easy, out-patient procedure which may be needed 6-12 months following surgery. The implants (bone plates) do not cause any problems in some patients and may not need to be removed. Recovery following removal of the bone plate is minimal (4 weeks to allow the incision to heal and screw holes to fill in) with fewer risks of complications.